

## **Brown warts**

### **Also known as**

Seborrheic Keratosis, Age warts, senile warts

### **What are seborrheic keratoses?**

Seborrheic keratoses are benign “wart-like” growths on the skin. Seborrheic keratoses affect all racial groups and most commonly appear after the age of 40 years although some people may develop them earlier.

They are often considered unsightly in appearance.

### **What causes seborrheic keratoses?**

Seborrheic keratoses are benign (non-cancerous) growths of the skin.

The exact cause of seborrheic keratoses is unknown although they do tend to occur more frequently with increasing age. Their appearance is thought to be part of the normal ageing process of the skin.

### **What do seborrheic keratoses look like?**

Seborrheic keratoses are “wart-like” growths. They are usually raised above the surface of the skin and appear to be stuck on like “barnacles”.

Seborrheic keratoses occur in varying shades of brown from light to very dark brown. The surface of seborrheic keratoses usually feels rough and warty. They can sometimes be dry and scaly and become itchy or irritated.

Most people will have more than one seborrheic keratosis and some people may have hundreds scattered over the body. They can appear anywhere on the body including the scalp, face and genitals. In a condition called dermatosis papulosis nigra hundreds of small seborrheic keratoses are present on the face and neck.

Sometimes seborrheic keratoses are flat and pale and can resemble freckles. Sometimes they can be large (up to many centimeters in size) and may resemble skin tumours.

They often appear quite suddenly.

### **What other problems may occur with seborrheic keratoses?**

In rare cases the sudden development of widespread, multiple (hundreds) seborrheic keratoses may be associated with an underlying malignant tumour.

### **How are seborrheic keratoses diagnosed?**

The diagnosis of a seborrheic keratosis is usually made by clinical examination.

Very dark lesions which have changed in some way or which are growing rapidly may need a biopsy to confirm the diagnosis and exclude the possibility of skin cancer.

Darker lesions that look irregular should be checked by a doctor to ensure that they are not melanoma.

**How are seborrheic keratosis treated?**

Seborrheic keratosis does not need any specific treatment.

Usually seborrheic keratosis are treated for cosmetic reasons or because they have become itchy or irritated. People affected with seborrheic keratosis often complain that they “catch” on things, bleed or become inflamed, especially if they are large and warty.

Seborrheic keratosis is usually treated by freezing them with liquid nitrogen (cryotherapy). They can also be removed surgically by lightly scraping (shave excision or curettage) or burning the lesions (electrocautery). Laser resurfacing (ablative laser) and strong medical-grade chemical peels may also be used.

All methods of removal have an increased risk of permanent colour change (lightening or darkening) or scarring of the skin. Removing seborrheic keratosis will not prevent new lesions from developing in the future.

There is no known effective prevention of these lesions.

**What is the prognosis or likely outcome of seborrheic keratosis?**

Seborrheic keratosis does not cause any major health problems. Some people affected by this condition may be troubled by the cosmetic appearance of the lesions. Lesions can be traumatized accidentally by scratching and may bleed or become itchy or inflamed.